

# **Potassium Starvation in Yeast Triggers Changes in the Expression of Genes Related to Different Metabolic and Biosynthetic Pathways**

Aby Mesfin, Jonar Cowan, David Ramirez, Mihir Samdarshi,  
Christina Dominguez  
Loyola Marymount University  
BIOL 367-01  
December 10, 2019

# Outline

- Potassium crucial to cell
- Raw Data Required Manual Input
- Multiple significant changes found in genes
- Eight Significant Gene Clusters Observed
- Profile 39 was most significant
- Top Transcription Factor Genes Serve Importance in Cell
- 12 Gene Network
- Database
- Network Discovered

# Outline

- **Potassium crucial to cell**
- Raw Data Required Manual Input
- Multiple significant changes found in genes
- Eight Significant Gene Clusters Observed
- Profile 39 was most significant
- Top Transcription Factor Genes Serve Importance in Cell
- 12 Gene Network
- Database
- Network Discovered

# Potassium is extremely important to normal cell function

- Potassium is the most abundant cation
- Without potassium balance life threatening conditions might occur
- Potassium homeostasis is required to regulate the cell
- Potassium helps to establish membrane potential
- Potassium limits growth when ammonium is the nitrogen source
- Ammonium toxicity is increased under high-potassium conditions
  - Caused by over-expression of ammonium transporters

# Potassium is extremely important to normal cell function

- “The Short-Term Response of Yeast to Potassium Starvation” by Barreto, L., Canadell, D., Valverde-Saubí, D., Casamayor, A., & Ariño, J.
  - Study looked at potassium starvation in yeast cells for 10, 20, 40, 60, and 120 minutes in replicates of four
    - Found changes in gene expression impacted by potassium starvation
      - Oxidative Stress
      - Methionine/Cysteine Biosynthesis
      - Cyclin Levels
      - Septin Rings
      - Retrograde Pathway
      - Methylglyoxal production
      - Trehalose metabolism
  - No studies that directly correlated to work on potassium starvation prior to this study


# Potassium is extremely important to normal cell function

- GRNmap and GRNsight help visualize the relationship between transcription factors during potassium starvation
- MS Access database will be used for present and future use of understanding and exploring new information from microarray data

# Outline

- Potassium crucial to cell
- **Raw Data Required Manual Input**
- Multiple significant changes found in genes
- Eight Significant Gene Clusters Observed
- Profile 39 was most significant
- Top Transcription Factor Genes Serve Importance in Cell
- 12 Gene Network
- Database
- Network Discovered

# Complications with the Raw Data





# Outline

- Potassium crucial to cell
- Raw Data Required Manual Input
- **Multiple significant changes found in genes**
- Eight Significant Gene Clusters Observed
- Profile 39 was most significant
- Top Transcription Factor Genes Serve Importance in Cell
- 12 Gene Network
- Database
- Network Discovered

# More than 5% of the Genes Had A Significant Change in Gene Expression

<b>ANOVA</b>	<b>Percent of Genes</b>
<b>p &lt; 0.05</b>	2,985 (71.9 %)
<b>p &lt; 0.01</b>	2,403 (57.9 %)
<b>p &lt; 0.001</b>	1,703 (41 %)
<b>p &lt; 0.0001</b>	1,198 (28.9 %)
<b>B &amp; H p &lt; 0.05</b>	2,839 (68.4 %)
<b>Bonferroni p &lt; 0.05</b>	776 (18.7 %)

# Outline


- Potassium crucial to cell
- Raw Data Required Manual Input
- Multiple Significant Change Found in Genes
- **Eight Significant Gene Clusters Observed**
- Profile 39 was most significant
- Top Transcription Factor Genes Serve Importance in Cell
- 12 Gene Network
- Database
- Network Discovered

# Eight Significant Gene Clusters Observed



Profile 39 was the most Significant Cluster


Profile 39 was used for the rest of the study



# Outline

- Potassium crucial to cell
- Raw Data Required Manual Input
- Multiple Significant Change Found in Genes
- Eight Significant Gene Clusters Observed
- **Profile 39 was most significant**
- Top Transcription Factor Genes Serve Importance in Cell
- 12 Gene Network
- Database
- Network Discovered

# Gene Plot of Profile 39 Displayed Significant Changes at Every Time-Point



# Outline

- Potassium crucial to cell
- Raw Data Required Manual Input
- Multiple Significant Change Found in Genes
- Eight Significant Gene Clusters Observed
- Profile 39 was most significant
- **Top Transcription Factor Genes Serve Importance in Cell**
- 12 Gene Network
- Database
- Network Discovered

# F/C in Genes Associated with Cellular Functions

Category Name	#Genes Category	#Genes Assigned	#Genes Expected	#Genes Enriched	p-value	Corrected p-value	Fold
cysteine biosynthetic process	10	9	1.2	7.8	3.10E-08	<0.001	7.8
cellular amino acid biosynthetic process	67	24	7.8	16.2	1.40E-07	<0.001	3.1
methionine biosynthetic process	26	13	3	10	1.40E-06	<0.001	4.3
hydrogen sulfide biosynthetic process	6	6	0.7	5.3	2.30E-06	<0.001	8.6
molecular function	694	114	80.4	33.6	6.20E-06	<0.001	1.4
oxidation-reduction process	211	45	24.5	20.5	1.80E-05	0.004	1.8



# Top 16 Significant Transcription Factor Genes From Yeastract

RPN4, PDR1, PDR3, HSF1  
Removed

- Fold change data for these genes was not measured in the study
- Empty query results


Transcription Factor	% in <i>s. cerevisiae</i>
Gat1p	0.6079
Gcn4p	0.9878
Met4p	0.2827
Rpn4p	0.9635
Yrr1p	0.535
Pdr1p	0.9149
Pdr3p	0.9726
Hsf1p	0.5015

Transcription Factor	% in <i>s. cerevisiae</i>
Bas1p	0.7964
Hac1p	0.3951
Mga2p	0.3587
Rlm1p	0.2766
Sok2p	0.5927
Fhl1p	0.5137
Arr1p	0.3313
Gis1p	0.2462


# Outline

- Potassium crucial to cell
- Raw Data Required Manual Input
- Multiple Significant Change Found in Genes
- Eight Significant Gene Clusters Observed
- Profile 39 was most significant
- Top Transcription Factor Genes Serve Importance in Cell
- **12 Gene Network**
- Database
- Network Discovered

# Regulatory Network Established Relationship between 12 Genes



# Regulatory Network Established Relationship between 12 Genes




# Outline

- Potassium crucial to cell
- Raw Data Required Manual Input
- Multiple Significant Change Found in Genes
- Eight Significant Gene Clusters Observed
- Profile 39 was most significant
- Top Transcription Factor Genes Serve Importance in Cell
- 12 Gene Network
- **Database**
- Network Discovered

# Database of DNA microarray data was created in Microsoft Access using existing database

## Windows Virtual Machine - Microsoft Azure



## Existing database utilized as starting point

### All Access Objects

Search...

#### Tables

- dcin5\_log2\_expression
- degradation\_rates
- dglin3\_log2\_expression
- dhap4\_log2\_expression
- dzap1\_log2\_expression
- production\_rates
- wt\_log2\_expression




### All Access Objects

Search...

#### Tables

- Barreto\_2012\_wt\_log2\_expression
- Dahlquist\_2018\_dcin5\_log2\_expression
- Dahlquist\_2018\_dglin3\_log2\_expression
- Dahlquist\_2018\_dhap4\_log2\_expression
- Dahlquist\_2018\_dzap1\_log2\_expression
- Dahlquist\_2018\_wt\_log2\_expression
- degradation\_rates
- Kitagawa\_2002\_wt\_log2\_expression
- production\_rates
- references
- samples\_to\_data
- Thorsen\_2007\_wt\_log2\_expression

# Diagram describes MS Access database schema



# Two new tables were added containing data source metadata

`samples_to_data` and `references` Tables

	Field Name	Data Type
PK	Sample ID	Number
	PubMed ID	Number
	Control Yeast Strain	Short Text
	Treatment Yeast Strain	Short Text
	Control	Short Text
	Treatment	Short Text
	Concentration Value	Number
	Concentration Unit	Short Text
	Time Value	Number
	Time Units	Short Text
	Number of Replicates	Number
	Table	Short Text

	Field Name	Data Type
PK	Pubmed ID	Number
	Authors	Short Text
	Publication Year	Number
	Title	Short Text
	DOI	Short Text
	NCBI GEO ID	Short Text



# Field and table names were standardized across gene expression data tables

## Gene Expression Data Table Name

Format: [Author 1 Surname] [Year] [wt or mutant]\_log2\_expression

Example: Barreto\_2012\_wt\_log2\_expression

## Gene Expression Data Field Name (in Table)

Format: [Yeast Strain]\_LogFC\_t[Time Point]-[Replicate Number]

Example: Barreto\_2012\_wt\_log2\_expression

# Outline

- Potassium crucial to cell
- Raw Data Required Manual Input
- Multiple Significant Change Found in Genes
- Eight Significant Gene Clusters Observed
- Profile 39 was most significant
- Top Transcription Factor Genes Serve Importance in Cell
- 12 Gene Network
- Database
- **Network Discovered**

# A Gene Regulatory Network Was Induced in the Yeast by Potassium Starvation

- Potassium is proven to play crucial role in cell function
- Raw data needed multiple edits including deletion of aliases
- More than 5% of the genes had a significant change in gene expression
- Eight significant gene clusters
  - Profile 39 → most significant
  - Significant time points for every change
- Important relationship observed between 12 genes
  - 4/16 original top transcription factors had no data
  - Genes related to roles in cellular function

# Genes Highlighted in Our Regulatory Network Are Absent in the Article

- Genes in the Article are associated with:
  - Oxidative Stress
  - Methionine/Cysteine Biosynthesis
  - Cyclin Levels
  - Septin Rings
  - Retrograde Pathway
  - Methylglyoxal production
  - Trehalose metabolism
- Our Network highlights genes associated with:
  - Amino Acid Biosynthesis
  - Methionine/Cysteine Biosynthesis
  - Hydrogen Sulfide Biosynthesis
  - Molecular Function
  - Oxidation-Reduction Processes

# Network and GO Definitions Are Substantiated by Previous Studies

- Bas1 is a regulator of the Histidine Pathway and Purine Biosynthetic Pathway (Daignan-Fornier, et al., 1920)
- Fhl1 functions as a suppressor of mutations in RNA Polymerase III (Hermann-Le Denmat, et al., 1994)
- Gat1 and Fhl1 involved in similar metabolic pathways (Bandhakavi, 2008)
- Gcn4 regulates gene expression during amino acid starvation in yeast (Natarajan, et al., 2010)
  - Activator of Gat1, Met4, Bas1
- Sok2 and Gis1 cooperate in regulating the promoter of TPK1 (Pautasso, Rossi, 2014)

# Network and GO Definitions Are Substantiated by Previous Studies

- Met4 related to sulfur metabolism and binding factor both needed for transcriptional activation (Thomas et al., 2004)
- Hac1 and Mga2 found to be related in cellular function but exact relationship unknown (Covino et al. 2018)
- Rlm1 delays the transition from G1 to S in cell cycle (Piccirillo et al., 2017)
- Yrr1 responds to oxidative stress (Nadai et al., 2016)
- Gls1 induces other metabolic genes in catabolic pathways (Wang et al., 2011)

# Discovering More About Network

- Finding the 4 missing gene expressions
- How the 12 significant genes affect:
  - Other cations and the effects on the cell
  - Gene expression
  - Cellular Function in relation to homeostasis and membrane potential

# Acknowledgments

Thank You  
Dr. Dahlquist  
LMU Biology Department  
BIOL 367 Classmates



# References

- Bandhakavi, S., Xie, H., O'Callaghan, B., Sakurai, H., Kim, D. H., & Griffin, T. J. (2008). Hsf1 activation inhibits rapamycin resistance and TOR signaling in yeast revealed by combined proteomic and genetic analysis. *PloS one*, 3(2), e1598.
- Barreto, L., Canadell, D., Valverde-Saubí, D., Casamayor, A., & Ariño, J. (2012). The short-term response of yeast to potassium starvation. *Environmental microbiology*, 14(11), 3026-3042. DOI: 10.1111/j.1462-2920.2012.02887.x
- Covino, R., Hummer, G., & Ernst, R. (2018). Integrated functions of membrane property sensors and a hidden side of the unfolded protein response. *Molecular cell*, 71(3), 458-467.
- Daignan-Fornier, B., & Fink, G. R. (1992). Coregulation of purine and histidine biosynthesis by the transcriptional activators BAS1 and BAS2. *Proceedings of the National Academy of Sciences*, 89(15), 6746-6750.
- Flick, K., Ouni, I., Wohlschlegel, J. A., Capati, C., McDonald, W. H., Yates 3rd, J. R., & Kaiser, P. (2004). Proteolysis-independent regulation of the transcription factor Met4 by a single Lys 48-linked ubiquitin chain. *Nature Cell Biology*, 6(7), 634.
- Guerfal, M., Ryckaert, S., Jacobs, P. P., Ameloot, P., Van Craenenbroeck, K., Derycke, R., & Callewaert, N. (2010). The HAC1 gene from *Pichia pastoris*: characterization and effect of its overexpression on the production of secreted, surface displayed and membrane proteins. *Microbial cell factories*, 9, 49. doi:10.1186/1475-2859-9-49
- Hermann-Le Denmat, S., Werner, M., Sentenac, A., & Thuriaux, P. (1994). Suppression of yeast RNA polymerase III mutations by FHL1, a gene coding for a fork head protein involved in rRNA processing. *Molecular and cellular biology*, 14(5), 2905-2913.

# References

- Nadai, C., Treu, L., Campanaro, S., Giacomini, A., & Corich, V. (2016). Different mechanisms of resistance modulate sulfite tolerance in wine yeasts. *Applied microbiology and biotechnology*, *100*(2), 797-813.
- Natarajan, K., Meyer, M. R., Jackson, B. M., Slade, D., Roberts, C., Hinnebusch, A. G., & Marton, M. J. (2001). Transcriptional profiling shows that Gcn4p is a master regulator of gene expression during amino acid starvation in yeast. *Molecular and cellular biology*, *21*(13), 4347-4368.
- Pautasso, C., & Rossi, S. (2014). Transcriptional regulation of the protein kinase A subunits in *Saccharomyces cerevisiae*: Autoregulatory role of the kinase A activity. *Biochimica Et Biophysica Acta (BBA)-Gene Regulatory Mechanisms*, *1839*(4), 275-287.
- Piccirillo, S., Neog, D., Spade, D., Van Horn, J. D., Tiede-Lewis, L. M., Dallas, S. L., ... Honigberg, S. M. (2017). Shrinking Daughters: Rlm1-Dependent G1/S Checkpoint Maintains *Saccharomyces cerevisiae* Daughter Cell Size and Viability. *Genetics*, *206*(4), 1923–1938. doi:10.1534/genetics.117.204206
- Wang, R., Dillon, C. P., Shi, L. Z., Milasta, S., Carter, R., Finkelstein, D., ... & Green, D. R. (2011). The transcription factor Myc controls metabolic reprogramming upon T lymphocyte activation. *Immunity*, *35*(6), 871-882.
- Zhang, S., Skalsky, Y., & Garfinkel, D. J. (1999). MGA2 or SPT23 is required for transcription of the  $\Delta 9$  fatty acid desaturase gene, OLE1, and nuclear membrane integrity in *Saccharomyces cerevisiae*. *Genetics*, *151*(2), 473-483.